

BELGISCHE BODEMKUNDIGE VERENIGING - SOCIÉTÉ BELGE DE PÉDOLOGIE

BELGISCHE BODENKUNDLICHE GESELLSCHAFT - BELGIAN SOIL SCIENCE SOCIETY

PÉDOLOGIE - THEMATA

1998 - N° 5

Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen, RUG,
Coupure 653, B-9000 Gent, België

Comparing pedotransfer functions to estimate soil bulk density in northern Belgium

BOUCNEAU G., VAN MEIRVENNE M. & HOFMAN G.

University Gent, Dept. Soil Management, Coupure 653, 9000 Gent, Belgium

INTRODUCTION

Soil bulk density (D) is required for estimating soil water retention characteristics and is an often required input parameter for water, sediment and nutrient transport models. Direct measurements of D are based on undisturbed samples and so they are labor-intensive and time-consuming. Consequently it is not routinely analysed. Much effort has been devoted to develop pedotransfer functions (PTF) (Bouma, 1989) which relate D with available basic soil properties.

D depends on the specific gravity of the soil particles and their arrangement. Since the specific gravity of soil minerals is more or less constant, soil structure is chiefly responsible for fluctuations of bulk density. The structural properties, in turn, are determined largely by the interacting influences of the soil particles and the degree of aggregation for which organic matter is responsible. As a consequence D is found to be related to soil texture and soil organic matter (Saini, 1966 and Adams, 1973).

Mostly PTF were developed for particular soil groups and so these have a limited applicability (Van Wambeke, 1974 ; Alexander, 1980 ; Mbagwu *et al.*, 1983 and Scott & Wood, 1989). Brakensiek *et al.* (1980) and McCuen *et al.* (1981) published class PTF with average D values for the USDA soil texture classes. PTF which are relevant for a wide range of soils were developed by Adams (1973) ; Rawls (1983) and Manrique & Jones (1991). Van Hove (1969) and Van Ruymbeke *et al.* (1991) developed class PTF for northern Belgium. Until now, no continuous PTF for D of soils in Northern Belgium were developed. The aim of

this study was to compare PTF from literature to estimate soil bulk density in northern Belgium.

MATERIALS AND METHODS

Class pedotransfer functions

If no direct measurements are made, D of soils in northern Belgium is estimated using the two class PTF published in Van Hove (1969) and Van Ruymbeke *et al.* (1991). Van Ruymbeke *et al.* (1991) estimated D as being 1.45 g cm^{-3} for topsoil horizons (rich in organic matter) and 1.50 g cm^{-3} for underlying horizons (low in organic matter). Van Hove (1969) refined this by relating D of topsoil horizons of northern Belgium to the classes of the Belgian soil texture triangle (Figure 1).




Figure 1 D of topsoil horizons in function of the classes on the Belgian soil texture triangle (Van Hove, 1969).

Continuous pedotransfer functions

Adams (1973) predicts D using the equation:

$$D = \frac{100}{\frac{OM}{D_{OM}} + \frac{100-OM}{D_M}} \quad \text{Eq. 1}$$

with OM the organic matter content, D_{OM} the bulk density of the organic matter which is set to 0.224 g cm^{-3} and D_M the bulk density of the mineral fraction. Rawls (1983) published a contour map of D_M as a function of sand and clay content (Figure 2).

Manrique & Jones (1991) predicted D as a function of the square root of organic carbon (C):

$$D = 1.660 - 0.318\sqrt{C} \quad \text{Eq. 2}$$

They also grouped soils according to the USDA soil orders and determined a functional relation for each order. We translated the major Belgian soil series to USDA soil orders: Alfisols (Aba, Aca, Lba, Lca, Lda, sLba, Ldc, Sbc,), Entisols (Abp, sEdp, Udp), Inceptisols (Sbf, Sdm) and Podisols (Zdg, Zcg):

$$D = 1.704 - 0.005Cl - 0.127C \quad \text{Eq. 3}$$

for Alfisols

$$D = 1.861 - 0.452\sqrt{C} \quad \text{Eq. 4}$$

for Entisols

$$D = 1.720 - 0.403\sqrt{C} \quad \text{Eq. 5}$$

for Inceptisols

$$D = 1.587 - 0.259\sqrt{C} \quad \text{Eq. 6}$$

for Spodosols

with Cl the percentage of clay.

Validation data

The dataset published by Vereecken (1988) was used to compare these PTF. The dataset contains information on the soil textural fractions, organic matter and bulk density for

the horizons (182) of 40 profiles representing the major soil series in northern Belgium.




Figure 2 Contour map of D_M (g cm^{-3}) (Rawls, 1983)

The validation indices used to compare the PTF were the mean prediction error (MPE), the mean and the standard deviation of the square prediction error ($MSPE$ and $SDSPE$ respectively) and the Pearson correlation coefficient (r), defined as:

$$MPE = \frac{1}{n} \sum_{i=1}^n (D_i - \hat{D}_i) \quad \text{Eq. 7}$$

$$MSPE = \frac{1}{n} \sum_{i=1}^n (D_i - \hat{D}_i)^2 \quad \text{Eq. 8}$$

$$SDSPE = \sqrt{\frac{\sum_{i=1}^n ((D_i - \hat{D}_i)^2 - MSPE)^2}{n-1}} \quad \text{Eq. 9}$$

$$r = \frac{\text{Cov}(D_i, \hat{D}_i)}{\sqrt{\text{Var}(D_i)\text{Var}(\hat{D}_i)}} \quad \text{Eq. 10}$$

In these equations D_i is the observed soil bulk density, \hat{D}_i is the soil bulk density predicted

by the PTF and n is the number of observations (182). The prediction should be unbiased, so MPE should approach zero. $MSPE$ is a measure of the average precision, $SDSPE$ is a measure of the variation in precision and both should be as small as possible. The Pearson correlation coefficient is a measure of the strength of similarity between observations and predictions.

RESULTS

Table 1 presents the validation values for the five PTF. For the class PTF (Van Hove, 1969 and Van Ruymbeke *et al.*, 1991) MPE s are close to zero and $MSPE$ s are low. However the relationship between predictions and observations is poor ($r < 0.3$) (Figure 3a-b) and also the $SDSPE$ s have intermediate values.

Table 1 Performance of PTF for D

PTF	MPE $g\ cm^{-3}$	$MSPE$ $g^2\ cm^{-6}$	$SDSPE$ $g^2\ cm^{-6}$	r
Van Ruymbeke <i>et al.</i> (1991)	0.007	0.0171	0.0407	0.17
Van Hove (1969)	0.025	0.0177	0.0410	0.25
Rawls (1983)	0.173	0.0501	0.0555	0.50
General PTF of Manrique & Jones (1991)	0.117	0.0305	0.0417	0.58
PTF based on the USDA orders (Manrique & Jones, 1991)	0.015	0.0184	0.0309	0.61




Figure 3 Estimated versus observed D for different PTF: (a) Van Ruymbeke *et al.* (1991), (b) Van Hove (1969), (c) Rawls (1983), (d) general PTF of Manrique & Jones (1991), (e) PTF based on the USDA orders of Manrique & Jones (1991).

For Manrique & Jones (1991) and Rawls (1983) the high *MSPEs* and *SDSPEs* indicate a poor accuracy, induced by an important systematic underestimation as indicated by the *MPE* and as can be visually observed in Figure 3c-d. However $r's \geq 0.50$ indicate a stronger relationship between predictions and observations than for the two class PTF.

The PTF based on the USDA orders (Manrique & Jones, 1991) score the best for the Pearson correlation coefficient (Figure 3e) and *SDSPE*. The *MPE* and *MSPE* approach the minimal values obtained by the class PTF. This study also points out that the evaluation of predictive methods should be based on several validation indices simultaneously.

CONCLUSIONS

For soils in northern Belgium predictions of bulk density are best performed by the PTF on the basis of the USDA orders (Manrique & Jones, 1991). If only the texture class is known, the class PTF of Van Hove (1969) is the best alternative.

REFERENCES

- Adams, W. A. 1973. The effect of organic matter on the bulk density and true densities of some uncultivated podzolic soils. *The Journal of Soil Science* 24:10-17.
- Alexander, E. B. 1980. Bulk densities of Californian soils in relation to other soil properties. *Soil Sci. Soc. Am. J.* 44:689-692.
- Bouma, J. 1989. Using soil survey data for quantitative land evaluation. *Advances in Soil Science* vol. 9 (ed. Stewart), Springer-Verlag, pp. 177-213.
- Brakensiek, D. L. ; Engleman, R. L. & Rawls, W. J. 1980. Variation within soil texture classes of soil water parameters. *Trans. Am. Soc. Agric. Eng.* 24:335-339.
- Manrique, L. A. & Jones C. A. 1991. Bulk density of soils in relation to soil physical and chemical properties. *Soil Sci. Soc. Am. J.* 55:476-481.
- Mbagwu, J. ; Lal, R. & Scott, T. W. 1983. Physical properties of three soils in southern Nigeria. *Soil Science* 136: 123-125.
- McCuen, R. H. ; Rawls, W. J. & Brakensiek, D.L. 1981. Statistical analyses of the Brooks and Corey and Green-Ampt parameters across soil texture. *Water Resour. Res.* 17:1005-1013.
- Rawls, W. J. 1983. Estimating soil bulk density from particle size analysis and organic matter content, *Soil Science* 135:123-125.
- Saini, G. R. 1966. Organic matter as a measure of bulk density of soil. *Nature* 210:295-1296.
- Scott, H. D. & Wood, L. S. 1989. Impact of crop production on the physical status of a Typic Albaqualf. *Soil Sci. Soc. Am. J.* 53:1782-17789.
- Van Hove, J. 1969. Variatie van het gehalte aan organisch materiaal en van de C/N-verhouding in de oppervlakte-horizonten van de bodems van Laag- en Midden-België. Aggregaat voor het Hoger Onderwijs. Universiteit Gent, 291 p.
- Van Ruymbeke, M. ; Hofman, G. ; Demyttemaere, P. ; Verstegen, P. & Van Meirvenne, M. 1991. Organische stof problematiek en kwantificatie van de beschikbare voedingselementen in de Westvlaamse groentestreek. I.W.O.N.L. rapport 1ste biënnale, Conventie D1/4 - 12489/5332A, 61 p.
- Vereecken, H. 1988. Pedotransfer functions for the generation of hydraulic properties for Belgian soils. Ph.D.thesis, Catholic University of Leuven, 254 p
- Van Wambeke, A. 1974. Management properties of Ferralsols. *Soils Bull.* 23, FAO, Rome, Italy.